

REVIEW

Surveying the epigenomic landscape, one base at a time

Gabriel E Zentner¹ and Steven Henikoff^{1,2,*}

Abstract

Epigenomics, the determination of epigenetic landscapes on a genome-wide scale, has progressed at an astounding rate over the past decade. Recent technological developments have enabled base-pair resolution of various epigenomic features, leading to new insights into epigenetic regulation.

Keywords epigenomics, chromatin immunoprecipitation, ChIP-exo, micrococcal nuclease mapping, deoxyribonuclease I mapping

The epigenome

Eukaryotic genomes are packaged into chromatin, consisting mostly of nucleosomes composed of approximately 147 bp of DNA wrapped around basic histone octamers [1]. Nucleosomes package DNA approximately 10,000-fold to form metaphase chromosomes, and so are essential for faithful segregation of sister genomes at mitosis. As nucleosomes occupy approximately 70% of the chromatin landscape during interphase, they must be mobilized during processes that require access to DNA, such as replication, transcription, repair, and binding by regulatory proteins. The occupancy, positioning, and composition of nucleosomes, as well as chemical modifications of histones and DNA, form a complex landscape superimposed on the genome: the epigenome [2]. Whereas the genome sequences of many organisms are now essentially complete [3], inquiry into their epigenomes is grossly incomplete due to the complexity and dynamics of the individual epigenomic constituents.

As in prokaryotes, sequence-specific DNA-binding proteins stand at the top of the eukaryotic transcriptional regulatory hierarchy, and differential expression of

transcription factors (TFs) results in cell type-specific differences. Most other key chromatin components are found in all cells of an organism, and dynamically change their distribution as a result of TF binding. The incorporation of histone variants [4] and the covalent modification of histone tails [5] help to mediate the inheritance of expression states of a gene by regulating the accessibility of DNA. Additionally, hundreds of chromatin-associated proteins, including ATP-dependent chromatin remodelers [6] and histone modifying enzymes [5], interact with chromatin to modulate its structure. Notably, mutations in nucleosome remodelers and in the histone constituents of chromatin have been implicated in human developmental disorders and cancer [6,7]. Thus, high-resolution genomic analysis of chromatin structure and the proteins that influence it is a major focus of biological technology development to study both basic cellular processes and the pathogenesis of human disease.

Many methods have been put forth with which to probe various aspects of the epigenome (Table 1), but until recently, the resolution of genome-wide methods for epigenome characterization, such as ChIP-chip [8] and MeDIP [9], was on the order of hundreds of base pairs, with the use of hybridization-based read-out technologies and chromatin preparation protocols based on random fragmentation. However, with the advent of massively parallel short-read DNA sequencing and its potential for single base-pair resolution, there has been a renaissance of interest in traditional methods for chromatin characterization, including the use of bisulfite sequencing for mapping DNA methylation [10] and the use of non-specific nucleases, including micrococcal nuclease (MNase) [11], deoxyribonuclease I (DNase I) [12] and exonuclease [13] (Table 1). Here, we focus on recently developed strategies for characterizing nucleosomes, TFs and chromatin-associated proteins at base-pair resolution, and we discuss prospects for full epigenome characterization.

Technologies for base-pair resolution epigenomic mapping

Several recent studies have introduced methods for analyzing various protein components of the epigenome

*Correspondence: steveh@fhrc.org

¹Basic Sciences Division, Fred Hutchinson Cancer Research Center, 1100 Fairview Ave N., Seattle, WA 98109, USA

²Howard Hughes Medical Institute, Fred Hutchinson Cancer Research Center, 1100 Fairview Ave N., Seattle, WA 98109, USA

Table 1. Strategies for epigenome mapping

Method	Strategy	Features	Base-pair resolution possible?	Reference
MNase digestion	Digests non-occluded DNA	Maps nucleosomes, other DNA-occluding particles	Yes [20,23]	[11]
DNase I hypersensitivity	Digests non-occluded DNA and nucleosomes with 10 bp periodicity	Maps 'open' chromatin	Yes [30,44]	[12]
Salt fractionation	Extracts chromatin from intact nuclei with increasing salt concentrations	Maps 'active' and 'nuclear matrix' chromatin	Yes, in combination with MNase digestion	[45]
ChIP	Affinity purification of specific chromatin fragments	Maps protein binding sites	Yes (see ChIP-exo)	[46]
DamID	DNA marking by tethered Dam methyltransferase	<i>In vivo</i> marking of protein-DNA interactions	No	[47]
FAIRE	Differential solubility of 'open' and nucleosomal chromatin by sonication and phenol/chloroform extraction	Maps 'open' chromatin	Maybe	[38]
Sono-seq	Differential solubility of 'open' and nucleosomal chromatin by sonication and phenol/chloroform extraction	Maps 'open' chromatin	Maybe	[39]
CATCH-IT	Metabolic labeling of histones	Measures nucleosome turnover	Maybe	[48]
Genome-wide psoralen crosslinking	Treatment of DNA with the intercalating agent psoralen	Measures helical tension of DNA	No	[49]
ChIP-exo	ChIP followed by exonuclease digestion of immunoprecipitated DNA	Removes non-occluded flanking DNA	Yes	[35]
Targeted chemical cleavage	Targeted chemical cleavage of DNA wrapped around modified nucleosomes	Maps nucleosome positions	Yes	[37]

Methods are listed in chronological order of their first published use in mapping epigenomic features to specific genomic locations. CATCH-IT, covalent attachment of tags to capture histones and identify turnover; ChIP, chromatin immunoprecipitation; ChIP-exo, chromatin immunoprecipitation with exonuclease digestion and high-throughput sequencing; DamID, DNA adenine methyltransferase identification; DNase I, deoxyribonuclease I; FAIRE, formaldehyde-assisted isolation of regulatory elements; MNase, micrococcal nuclease; Sono-seq, sonication of DNA and high-throughput sequencing.

at base-pair resolution while simultaneously addressing specific limitations of current epigenomic protocols. Below, we discuss the techniques upon which each of these high-resolution methods is based and how these new methods address the limitations of current epigenomic technologies.

MNase-seq

Digestion of chromatin with MNase has long been used to study chromatin structure in a low-throughput manner [14] and has more recently been combined with tiled microarray analysis (MNase-chip) or massively parallel DNA sequencing (MNase-seq) to study nucleosome positioning, occupancy, composition, and modification genome-wide [15]. MNase is a single-strand-specific secreted glycoprotein that is thought to cleave one strand of DNA as the helix breathes, then cleave the other strand to generate a double-strand break. MNase evidently 'nibbles' on the exposed DNA ends until it reaches an obstruction, such as a nucleosome. Though MNase has primarily been used to study nucleosomes, its mode of action suggests that it will be blocked by any obstruction along the DNA, such as a DNA-binding protein, allowing for the determination of genomic regions protected by non-histone proteins. By


combining MNase digestion with paired-end sequencing of protected DNA to determine precise fragment lengths, specific sizes of MNase-protected particles can be recovered with or without affinity purification and mapped. Indeed, we have used paired-end MNase-seq to map the distributions of both nucleosomes and paused RNA polymerase II in *Drosophila* cells [16]. Kent and colleagues [17] also used paired-end MNase-seq of native yeast chromatin to map the positions of both nucleosomes and sequence-specific TFs. Floer and colleagues [18] employed MNase digestion in conjunction with paired-end crosslinking chromatin immunoprecipitation (X-ChIP)-seq to identify binding sites for the RSC (remodels the structure of chromatin) complex, identifying partially unwrapped nucleosomes in the process. Importantly, these studies showed that DNA fragments as small as approximately 50 bp could be recovered after MNase digestion, suggesting applications for MNase-seq in epigenome mapping beyond nucleosome analysis.

A basic limitation of paired-end sequencing as a readout for MNase digestion, and epigenomic methods in general, is that standard short-read sequencing library preparation protocols are optimized for DNA fragments of nucleosomal size (approximately 150 bp) or larger and

involve size-selection of DNA [19], while regions of DNA protected by TFs are often up to an order of magnitude smaller. To circumvent this limitation, we introduced a modified library construction protocol to facilitate paired-end sequencing of DNA fragments as small as approximately 25 bp. By combining MNase digestion time points with mapping of a broad range of fragment sizes (approximately 25 to >200 bp), the distributions and dynamics of nucleosomes and non-histone proteins were analyzed [20]. Notably, subnucleosomal and nucleosomal particles can occupy the same genomic position within a population of cells, suggesting a highly dynamic interplay between nucleosomes and other chromatin-associated factors. As paired-end sequencing provides both fragment position and length, these two parameters can be displayed as a two-dimensional 'dot-plot'. The X-axis position of each dot represents the distance of the fragment midpoint to the center of a genomic feature such as a TF binding site (TFBS), and the Y-axis position represents its fragment length (Figure 1). The resulting

graph is referred to as a 'V-plot', because the minimal region of DNA protected is seen as the vertex of a 'V' corresponding to the fragment midpoint on the X-axis and its length on the Y-axis. Based on examination of V-plots for >100 TFs, the binding sites for TFs known to participate in nucleosome phasing, such as Abf1 and Reb1 [21,22], displayed well-positioned flanking nucleosomes and were flanked by subnucleosomal particles. V-plotting was also applied to ChIP data to show that the tripartite structure of the approximately 125-bp functional centromere sequence precisely corresponds to occupancy by a Cse4-containing centromeric nucleosome that is immediately flanked by particles corresponding to the Cbf1 TF and the kinetochore-specific Cbf3 complex [23].

MNase-seq with paired-end sequencing offers several advantages for epigenomic profiling. By mapping a wide range of fragment sizes, the genomic distributions of both nucleosomes and numerous non-histone proteins can be assessed using a single sequenced sample, making



the method especially cost-effective. The method does not require epitope tags or antibodies and is thus easily adapted to a range of cell types, particularly those for which affinity reagents are unavailable or impractical. No more than 25 cycles of sequencing per fragment end are needed to accurately map fragments onto genomes as large as that of *Drosophila* [24] and mouse (unpublished data), and the use of fewer cycles reduces both the cost and machine time for sequencing. Although MNase has a well-known AT-cleavage preference, in practice it causes only a minor mapping bias [25], which can be mitigated computationally if necessary [26]. A basic drawback of MNase-seq for mapping non-nucleosomal particles is that the identity of such particles cannot be formally established by this method alone, as multiple proteins may bind identical sequences. However, the recovery of non-nucleosomal particles from soluble native chromatin [20] suggests that this material is suitable for high-resolution ChIP-seq; indeed, it has been successfully applied to ChIP-seq mapping of paused RNA polymerase II in *Drosophila* [24]. The use of native chromatin for ChIP-seq (N-ChIP) may also offer solutions to issues associated with standard crosslinking ChIP protocols, such as epitope masking and protein-protein crosslinking due to formaldehyde treatment and the intrinsically low resolution of ChIP protocols employing sonication [27].

DNase-seq

DNase I is a non-specific endonuclease that has long been used for mapping sites of 'open' chromatin based on their hypersensitivity to cleavage [12]. Mapping of DNase I hypersensitivity with tiled microarrays (DNase-chip) or high-throughput sequencing (DNase-seq) has also been used to study the epigenome [28]. DNase I preferentially cleaves nucleosome-depleted genomic sites including regulatory elements such as promoters, enhancers, and insulators as well as TFBSs. DNase-seq identifies sites of DNase I digestion at base-pair resolution and offers an inverse approach to MNase-seq, as it infers the presence of DNA-occluding particles between hypersensitive sites whereas MNase maps the regions protected by such particles.

Hesselberth and colleagues [29] employed DNase-seq of yeast chromatin to map chromatin structure at computationally predicted binding sites for several TFs. Analysis of raw DNase-seq data revealed small regions of DNase protection within overall hypersensitive sites, likely indicative of TF binding. However, given that multiple proteins bind to identical sequences, it is necessary to integrate DNase-seq data with ChIP-seq data for definitive identification of the protein responsible for a particular DNase footprint. To this end, Boyle and colleagues [30] recently combined DNase-seq with TF ChIP-seq to precisely determine the DNA bound by

several TFs in human cells. Analysis of raw DNase-seq data revealed footprints of DNase resistance within larger hypersensitive sites, similar to the results of Hesselberth and colleagues [29]. DNase-seq was also central to the recent characterization of the human epigenome by the ENCODE consortium [31].

DNase-seq offers advantages to epigenomic analysis that are similar to MNase-seq in some respects. As it does not rely on antibodies or epitope tags, DNase-seq can query the genomic occupancy of numerous proteins in a single experiment and can be adapted to a range of cell types. However, given that multiple proteins can bind to identical sequences, integration of DNase-seq with ChIP-seq data is necessary to formally identify the protein responsible for a particular region of DNase protection. Mapping of nucleosome positioning with DNase-seq is also somewhat complicated, as DNase I cleaves nucleosomal DNA with 10 bp periodicity [32].

ChIP-exo

ChIP localizes proteins to specific sites on the genome and has become the most widely used epigenomic mapping technique in many fields of biological investigation. ChIP in combination with tiled microarray analysis (ChIP-ChIP) or high-throughput sequencing (ChIP-seq) has been extensively used to study the genomic distributions of hundreds of proteins [33]. While many important insights have been gained through ChIP-chip and ChIP-seq, there are limitations. Standard ChIP protocols employ sonication to fragment chromatin, which produces a heterogeneous mixture of fragments [34]. This issue is further compounded by size selection of 200 to 400 bp fragments during library preparation, a standard procedure in ChIP-seq protocols involving sonication [19]. Lastly, most ChIP-seq libraries are sequenced in single-end mode, wherein only one end of each DNA fragment is sequenced, and the resulting short sequence reads are computationally extended to approximate the size of each sequenced fragment. Taken together, these issues intrinsically limit the resolution of popular genome-wide ChIP methods.

To improve the resolution of ChIP-seq, Rhee and Pugh [35] introduced a technique called ChIP-exo. ChIP-exo involves performing a standard X-ChIP followed by λ exonuclease treatment. λ Exonuclease degrades DNA in a 5' to 3' manner and a protein crosslinked to DNA will block λ exonuclease digestion a specific number of bases 5' to the bound protein on each of the DNA strands, in effect creating a 5' barrier at a fixed distance from the protein past which exonuclease cannot digest and allowing sequences 3' of the barrier to remain intact. Following a specialized sequencing library preparation and single-end high-throughput sequencing, the 5' ends of the resulting sequence reads are mapped back to the

genome and precisely demarcate the 5' barriers created by the protein-DNA crosslinks with a high degree of precision and representing protein-bound locations as peak pairs, with one peak on either side of the bound protein. By precisely mapping the boundaries of exonuclease cleavage, ChIP-exo circumvents the limited resolution generally associated with single-end ChIP-seq.

ChIP-exo was applied to several yeast TFs, as well as to the human insulator-binding protein CTCF. Comparison of the yeast TF Reb1 ChIP-exo and ChIP-seq data revealed that ChIP-exo peaks displayed a standard deviation of 0.3 bp versus 24 bp for ChIP-seq peaks, a nearly 100-fold improvement in resolution [35]. The increased resolution of ChIP-exo revealed novel features regarding the modes of genomic binding by these factors. For instance, Reb1 displayed primary and secondary sites of occupancy. Secondary sites were defined as Reb1-occupied sites bound to a lesser extent than strongly occupied Reb1 sites within 100 bp. Notably, these primary-secondary Reb1 binding events were not resolved by standard ChIP-chip or ChIP-seq, indicating that ChIP-exo can resolve multiple crosslinking events within a single bound region. ChIP-exo analysis of other factors also revealed previously unidentified low-occupancy binding sites and refined and expanded the repertoire of sequence motifs associated with factor binding. For instance, CTCF occupancy was positively correlated with the presence of various sequence modules within a single consensus motif. CTCF binding sites with more modules tended to be more highly occupied, consistent with previous studies showing that CTCF uses various combinations of its 11 zinc fingers to bind distinct combinations of motif modules [36].

ChIP-exo addresses several limitations associated with conventional ChIP-seq. The precise mapping of nuclease protection boundaries allows for base-pair resolution determination of protein-bound sequences versus standard ChIP methodologies, which offer only an approximation of bound sequences. Additionally, unbound DNA contaminates ChIP samples, increasing background signal, which may lead to false positives in the case of highly enriched contaminating sequences and false negatives in the case of sites that are weakly bound by the protein of interest. Like MNase and DNase I, exonuclease treatment removes unbound DNA, greatly reducing the background of ChIP experiments (ChIP-exo signal-to-noise, 300- to 2,800-fold versus 7- and 80-fold for ChIP-chip and ChIP-seq, respectively), allowing for identification of low-occupancy binding sites and enabling in-depth analysis of relationships between DNA sequence and TF occupancy. Overall, ChIP-exo offers a base-pair resolution method by which to assess protein occupancy and further dissect the complex interplay between DNA sequence and TFs in genomic regulation

and should be readily applicable to systems with available ChIP reagents.

Adapting other epigenomic methods to single base-pair resolution mapping

MNase-seq, DNase-seq, and ChIP-exo, discussed above, are successful modifications of classical techniques for genome-wide analysis of epigenomic features. However, many other techniques have been used to map epigenomes (Table 1). One such technique is a novel targeted chemical cleavage approach that provides base-pair resolution mapping of nucleosome positions [37]. We therefore asked if other current techniques could be adapted for single base-pair resolution epigenome mapping.

Formaldehyde-assisted isolation of regulatory elements (FAIRE) [38] and Sono-seq [39] have been routinely used to map regions of 'open' chromatin. Both techniques rely on the fact that nucleosomes are much more readily crosslinked to DNA than are DNA-binding proteins when cells are treated with formaldehyde. While there are some differences in the FAIRE and Sono-seq protocols, they are based on the same principle. Cells are treated with formaldehyde to crosslink protein-DNA interactions and cells or isolated nuclei are sonicated to shear chromatin. Following sonication, samples are subjected to phenol-chloroform extraction. DNA not crosslinked to proteins ('open' chromatin) is recovered in the aqueous phase, while protein-DNA complexes are retained in the interface. DNA from the aqueous phase is then analyzed by microarray hybridization or high-throughput sequencing. However, as sonication produces a heterogeneous mixture of fragments and only non-protein-associated DNA is recovered, the precise positions of particles delimiting regions of 'open' chromatin cannot be obtained with these techniques. To map the precise positions of DNA-occluding particles using either the FAIRE or Sono-seq chromatin preparation protocol, the protein-DNA complexes contained in the insoluble fraction, which is normally discarded, could be purified and subjected to exonuclease digestion to generate DNA ends a uniform distance from each protein-DNA crosslink, as in ChIP-exo. High-throughput sequencing of exonuclease-digested chromatin would then reveal precise locations of DNA-protecting particles, and this approach could also be coupled to affinity purification to precisely localize specific factors.

Summary and future directions


While the development of technologies for base-pair resolution characterization of epigenomes is still in its early stages, important insights regarding chromatin organization have already been obtained with these

methods. ChIP-exo provides a method to precisely map the genomic binding of proteins in systems where ChIP reagents are readily available. MNase-seq allows for mapping of nucleosomes and non-histone proteins within a single sample and like DNase-seq is easily adapted to any system with a sequenced genome. In combination with ChIP-seq, MNase-seq and DNase-seq provide powerful methods for base-pair resolution identification of protein binding sites. These techniques are summarized schematically in Figure 2.

While epigenomic profiling is relatively straightforward in single-cell systems, it is more challenging in multicellular organisms, where different cell types are tightly interwoven in complex tissues. Indeed, ChIP-exo, MNase-seq, and DNase-seq have generally been performed either in yeast, which are unicellular, or cultured cells from other organisms, which are not necessarily reflective of the *in vivo* situation in the organism from which they were derived. To profile cell type-specific epigenomes at base-pair resolution, it will be necessary to combine the above technologies with methods for the isolation of specific cell types from a

complex milieu. One such method is fluorescence-activated cell sorting (FACS), involving purification of fluorescently labeled cells or nuclei. FACS has been used to isolate specific cell populations from mouse and human brain and mouse embryonic mesoderm for chromatin analysis [40,41]. Another technique, isolation of nuclei tagged in specific cell types (INTACT) has been used to isolate nuclei from individual cell types in *Arabidopsis*, *Caenorhabditis elegans*, and *Drosophila* for expression and preliminary chromatin profiling [42,43]. Combining these techniques with the various methods of base-pair resolution epigenome analysis detailed above should provide striking insights into the regulatory networks underlying specific cell identities.

As base-pair resolution epigenomic techniques are further developed and the cost of sequencing continues to decrease, genome-wide profiling of cell type-specific chromatin landscapes will become increasingly routine. The precise mapping of TFs, of nucleosomal features (positioning, occupancy, composition, and modification), and of ATP-dependent chromatin remodelers may provide the epigenomic equivalent of genome sequencing



projects, delineating the regulatory frameworks by which the various cell types within an organism use the same genome to generate distinct cellular identities.

Abbreviations

bp, base pair; ChIP, chromatin immunoprecipitation; ChIP-chip, chromatin immunoprecipitation with tiled microarray analysis; ChIP-seq, chromatin immunoprecipitation with high-throughput sequencing; ChIP-exo, chromatin immunoprecipitation with exonuclease digestion and high-throughput sequencing; DNase I, deoxyribonuclease I; DNase-chip, deoxyribonuclease I digestion with tiled microarray analysis; DNase-seq, deoxyribonuclease I digestion with high-throughput sequencing; FAIRE, formaldehyde-assisted isolation of regulatory elements; MNase, micrococcal nuclease; MNase-chip, micrococcal nuclease digestion with tiled microarray analysis; MNase-seq, micrococcal nuclease digestion with high-throughput sequencing; Sono-seq, sonication of DNA and high-throughput sequencing; TF, transcription factor; TFBS, transcription factor binding site; X-ChIP, chromatin immunoprecipitation with formaldehyde crosslinking.

Competing interests

The authors declare that they have no competing interests.

Acknowledgements

We thank John Latham, Paul Talbert, and Frank Pugh for comments on the manuscript. Work in our lab is supported by NIH grants 5U01 HG004274, U54 CA143862, and R01 ES020116 and by the Howard Hughes Medical Institute.

Published: 22 October 2012

References

- Luger K, Mader AW, Richmond RK, Sargent DF, Richmond TJ: **Crystal structure of the nucleosome core particle at 2.8 Å resolution.** *Nature* 1997, **389**:251-260.
- Bernstein BE, Meissner A, Lander ES: **The mammalian epigenome.** *Cell* 2007, **128**:669-681.
- Flicek P, Amode MR, Barrell D, Beal K, Brent S, Carvalho-Silva D, Clapham P, Coates G, Fairley S, Fitzgerald S, Gil L, Gordon L, Hendrix M, Hourlier T, Johnson N, Kähäri AK, Keefe D, Keenan S, Kinsella R, Komorowska M, Koscielny G, Kulesha E, Larsson P, Longden I, McLaren W, Muffato M, Overduin B, Pignatelli M, Pritchard B, Riat HS, et al.: **Ensembl 2012.** *Nucleic Acids Res* 2012, **40**:D84-D90.
- Talbert PB, Henikoff S: **Histone variants – ancient wrap artists of the epigenome.** *Nat Rev Mol Cell Biol* 2010, **11**:264-275.
- Bannister AJ, Kouzarides T: **Regulation of chromatin by histone modifications.** *Cell Res* 2011, **21**:381-395.
- Hargreaves DC, Crabtree GR: **ATP-dependent chromatin remodeling: genetics, genomics and mechanisms.** *Cell Res* 2011, **21**:396-420.
- Schwartzentruber J, Korshunov A, Liu XY, Jones DT, Pfaff E, Jacob K, Sturm D, Fontebasso AM, Quang DA, Tönjes M, Hovestadt V, Albrecht S, Kool M, Nantel A, Konermann C, Lindroth A, Jäger N, Rausch T, Ryzhova M, Korbel JO, Hielscher T, Hauser P, Garami M, Klekner A, Bogner L, Ebinger M, Schuhmann MU, Scheurlen W, Pekrun A, Frühwald MC, et al.: **Driver mutations in histone H3.3 and chromatin remodelling genes in paediatric glioblastoma.** *Nature* 2012, **482**:226-231.
- Iyer VR, Horak CE, Scafe CS, Botstein D, Snyder M, Brown PO: **Genomic binding sites of the yeast cell-cycle transcription factors SBF and MBF.** *Nature* 2001, **409**:533-538.
- Weber M, Davies JJ, Wittig D, Oakeley EJ, Haase M, Lam WL, Schubeler D: **Chromosome-wide and promoter-specific analyses identify sites of differential DNA methylation in normal and transformed human cells.** *Nat Genet* 2005, **37**:853-862.
- Harris RA, Wang T, Coarfa C, Nagarajan RP, Hong C, Downey SL, Johnson BE, Fouse SD, Delaney A, Zhao Y, Olshen A, Ballinger T, Zhou X, Forsberg KJ, Gu J, Echipare L, O'Geen H, Lister R, Pelizzola M, Xi Y, Epstein CB, Bernstein BE, Hawkins RD, Ren B, Chung WY, Gu H, Bock C, Gnirke A, Zhang MQ, Haussler D, et al.: **Comparison of sequencing-based methods to profile DNA methylation and identification of monoallelic epigenetic modifications.** *Nat Biotechnol* 2010, **28**:1097-1105.
- Reeves R, Jones A: **Genomic transcriptional activity and the structure of chromatin.** *Nature* 1976, **260**:495-500.
- Weintraub H, Groudine M: **Chromosomal subunits in active genes have an altered conformation.** *Science* 1976, **193**:848-856.
- Wu C: **An exonuclease protection assay reveals heat-shock element and TATA box DNA-binding proteins in crude nuclear extracts.** *Nature* 1985, **317**:84-87.
- Noll M: **Subunit structure of chromatin.** *Nature* 1974, **251**:249-251.
- Zhang Z, Pugh BF: **High-resolution genome-wide mapping of the primary structure of chromatin.** *Cell* 2011, **144**:175-186.
- Weber CM, Henikoff JG, Henikoff S: **H2A.Z nucleosomes enriched over active genes are homotypic.** *Nat Struct Mol Biol* 2010, **17**:1500-1507.
- Kent NA, Adams S, Moorhouse A, Paszkiewicz K: **Chromatin particle spectrum analysis: a method for comparative chromatin structure analysis using paired-end mode next-generation DNA sequencing.** *Nucleic Acids Res* 2011, **39**:e26.
- Floer M, Wang X, Prabhu V, Berrozpe G, Narayan S, Spagna D, Alvarez D, Kendall J, Krasnitz A, Stepansky A, Hicks J, Bryant GO, Ptashne M: **A RSC/ Nucleosome Complex Determines Chromatin Architecture and Facilitates Activator Binding.** *Cell* 2010, **141**:407-418.
- Schmidt D, Wilson MD, Spyrou C, Brown GD, Hadfield J, Odom DT: **ChIP-seq: Using high-throughput sequencing to discover protein-DNA interactions.** *Methods* 2009, **48**:240-248.
- Henikoff JG, Belsky JA, Krassovsky K, MacAlpine DM, Henikoff S: **Epigenome characterization at single base-pair resolution.** *Proc Natl Acad Sci U S A* 2011, **108**:18318-18323.
- Ganapathi M, Palumbo MJ, Ansari SA, He Q, Tsui K, Nislow C, Morse RH: **Extensive role of the general regulatory factors, Abf1 and Rap1, in determining genome-wide chromatin structure in budding yeast.** *Nucleic Acids Res* 2011, **39**:2032-2044.
- Hartley PD, Madhani HD: **Mechanisms that specify promoter nucleosome location and identity.** *Cell* 2009, **137**:445-458.
- Krassovsky K, Henikoff JG, Henikoff S: **Tripartite organization of centromeric chromatin in budding yeast.** *Proc Natl Acad Sci U S A* 2012, **109**:243-248.
- Teves SS, Henikoff S: **Heat shock reduces stalled RNA Polymerase II and nucleosome turnover genome-wide.** *Genes Dev* 2011, **25**:2387-2397.
- Allan J, Fraser RM, Owen-Hughes T, Keszenman-Pereyra D: **Micrococcal nuclease does not substantially bias nucleosome mapping.** *J Mol Biol* 2012, **417**:152-164.
- Albert I, Mavrich TN, Tomsho LP, Qi J, Zanton SJ, Schuster SC, Pugh BF: **Translational and rotational settings of H2A.Z nucleosomes across the *Saccharomyces cerevisiae* genome.** *Nature* 2007, **446**:572-576.
- O'Neill LP, Turner BM: **Immunoprecipitation of native chromatin: NChIP.** *Methods* 2003, **31**:76-82.
- Song L, Crawford GE: **DNase-seq: a high-resolution technique for mapping active gene regulatory elements across the genome from mammalian cells.** *Cold Spring Harb Protoc* 2010, **2010**:pdb.prot5384.
- Hesselberth JR, Chen X, Zhang Z, Sabo PJ, Sandstrom R, Reynolds AP, Thurman RE, Neph S, Kuehn MS, Noble WS, Fields S, Stamatoyannopoulos JA: **Global mapping of protein-DNA interactions in vivo by digital genomic footprinting.** *Nat Methods* 2009, **6**:283-289.
- Boyle AP, Song L, Lee B-K, London D, Keefe D, Birney E, Iyer VR, Crawford GE, Furey TS: **High-resolution genome-wide in vivo footprinting of diverse transcription factors in human cells.** *Genome Res* 2011, **21**:456-464.
- ENCODE Project Consortium: **An integrated encyclopedia of DNA elements in the human genome.** *Nature* 2012, **489**:57-74.
- Lutter LC: **Precise location of DNase I cutting sites in the nucleosome core determined by high resolution gel electrophoresis.** *Nucleic Acids Res* 1979, **6**:41-56.
- Raney BJ, Cline MS, Rosenbloom KR, Dreszer TR, Learned K, Barber GP, Meyer LR, Sloan CA, Malladi VS, Roskin KM, Suh BB, Hinrichs AS, Clawson H, Zweig AS, Kirkup V, Fujita PA, Rhead B, Smith KE, Pohl A, Kuhn RM, Karolchik D, Haussler D, Kent WJ: **ENCODE whole-genome data in the UCSC genome browser (2011 update).** *Nucleic Acids Res* 2011, **39**(Suppl 1):D871-D875.
- Schwartz YB, Kahn TG, Pirrotta V: **Characteristic low density and shear sensitivity of cross-linked chromatin containing polycomb complexes.** *Mol Cell Biol* 2005, **25**:432-439.
- Rhee Ho S, Pugh BF: **Comprehensive genome-wide protein-DNA interactions detected at single-nucleotide resolution.** *Cell* 2011, **147**:1408-1419.
- Ohlsson R, Lobanenkov V, Klenova E: **Does CTCF mediate between nuclear organization and gene expression?** *BioEssays* 2010, **32**:37-50.
- Brogaard K, Xi L, Wang J-P, Widom J: **A map of nucleosome positions in yeast at base-pair resolution.** *Nature* 2012, **486**:496-501.

38. Nagy PL, Cleary ML, Brown PO, Lieb JD: **Genomewide demarcation of RNA polymerase II transcription units revealed by physical fractionation of chromatin.** *Proc Natl Acad Sci U S A* 2003, **100**:6364-6369.
39. Auerbach RK, Euskirchen G, Rozowsky J, Lamarre-Vincent N, Moqtaderi Z, Lefrançois P, Struhl K, Gerstein M, Snyder M: **Mapping accessible chromatin regions using Sono-Seq.** *Proc Natl Acad Sci U S A* 2009, **106**:14926-14931.
40. Jiang Y, Matevosian A, Huang H-S, Straubhaar J, Akbarian S: **Isolation of neuronal chromatin from brain tissue.** *BMC Neurosci* 2008, **9**:42.
41. Takeuchi JK, Bruneau BG: **Directed transdifferentiation of mouse mesoderm to heart tissue by defined factors.** *Nature* 2009, **459**:708-711.
42. Deal RB, Henikoff S: **A simple method for gene expression and chromatin profiling of individual cell types within a tissue.** *Dev Cell* 2010, **18**:1030-1040.
43. Steiner FA, Talbert PB, Kasinathan S, Deal RB, Henikoff S: **Cell-type-specific nuclei purification from whole animals for genome-wide expression and chromatin profiling.** *Genome Res* 2012, **22**:766-777.
44. Boyle AP, Davis S, Shulha HP, Meltzer P, Margulies EH, Weng Z, Furey TS, Crawford GE: **High-resolution mapping and characterization of open chromatin across the genome.** *Cell* 2008, **132**:311-322.
45. Rocha E, Davie JR, van Holde KE, Weintraub H: **Differential salt fractionation of active and inactive genomic domains in chicken erythrocyte.** *J Biol Chem* 1984, **259**:8558-8563.
46. Gilmour DS, Lis JT: **In vivo interactions of RNA polymerase II with genes of *Drosophila melanogaster*.** *Mol Cell Biol* 1985, **5**:2009-2018.
47. van Steensel B, Henikoff S: **Identification of in vivo DNA targets of chromatin proteins using tethered Dam methyltransferase.** *Nat Biotechnol* 2000, **18**:424-428.
48. Deal RB, Henikoff JG, Henikoff S: **Genome-wide kinetics of nucleosome turnover determined by metabolic labeling of histones.** *Science* 2010, **328**:1161-1164.
49. Bermúdez I, García-Martínez J, Pérez-Ortín JE, Roca J: **A method for genome-wide analysis of DNA helical tension by means of psoralen-DNA photobinding.** *Nucleic Acids Res* 2010, **38**:e182.

doi:10.1186/gb-2012-13-10-250

Cite this article as: Zentner GE, Henikoff S: **Surveying the epigenomic landscape, one base at a time.** *Genome Biology* 2012, **13**:250.