PublisherInfo				
PublisherName		BioMed Central		
PublisherLocation		London		
PublisherImprintName		BioMed Central		

Schizophrenic NOTCH?

ArticleInfo		
ArticleID	:	4107
ArticleDOI	:	10.1186/gb-spotlight-20010605-01
ArticleCitationID	:	spotlight-20010605-01
ArticleSequenceNumber	:	178
ArticleCategory	:	Research news
ArticleFirstPage	:	1
ArticleLastPage	:	2
ArticleHistory	·	RegistrationDate : 2001–06–05 OnlineDate : 2001–06–05
ArticleCopyright	:	BioMed Central Ltd2001
ArticleGrants	:	
ArticleContext	:	130592211

Jonathan B Weitzman

Email: jonathanweitzman@hotmail.com

A recent report linked the human NOTCH4 locus with susceptibility to schizophrenia. In the June issue of Nature Genetics, two reports cast doubt on the NOTCH4-schizophrenia association. Sklar et al. conducted a large-scale study involving 519 parent-offspring trios in three independent families (compared to 80 trios in the original linkage study; Nature Genetics 2001, 28:126-128). Extensive association analyses failed to confirm the previous linkage results. McGinnis et al. analysed (CTG)n and (TAA)n microsatellite alleles in 300 Scottish schizophrenics and also failed to detect association (Nature Genetics 2001, 28:128-129). Thus, two large-scale studies employing transmission equilibrium test (TDT) analysis and case-control samples suggest that there is not a general association between NOTCH4 and schizophrenia.

References

- 1. The NOTCH4 locus is associated with susceptibility to schizophrenia.
- 2. Nature Genetics, [http://genetics.nature.com]
- 3. A powerful likelihood method for the analysis of linkage disequilibrium between trait loci and one or more polymorphic marker loci.