| PublisherInfo | | | | |
|----------------------|--------|----------------|--|--|
| PublisherName | | BioMed Central | | |
| PublisherLocation | | London | | |
| PublisherImprintName | \Box | BioMed Central | | |

Mighty splicing machine

| ArticleInfo | | |
|-----------------------|--|---|
| ArticleID | : | 4583 |
| ArticleDOI | | 10.1186/gb-spotlight-20020917-02 |
| ArticleCitationID | \Box | spotlight-20020917-02 |
| ArticleSequenceNumber | \Box | 249 |
| ArticleCategory | $\begin{bmatrix} \vdots \end{bmatrix}$ | Research news |
| ArticleFirstPage | \Box | 1 |
| ArticleLastPage | \Box | 2 |
| ArticleHistory | : | RegistrationDate : 2002–9–17 OnlineDate : 2002–9–17 |
| ArticleCopyright | | BioMed Central Ltd2002 |
| ArticleGrants | \Box | |
| ArticleContext | \Box | 130593311 |

Jonathan B Weitzman

Email: jonathanweitzman@hotmail.com

The spliceosome is a macromolecular machine, containing five small nuclear RNAs and a number of proteins, that is responsible for the excision of introns from pre-mRNA species. In the September 12 Nature, Zhou *et al.* report a complete proteomic analysis of all the polypeptide components of the spliceosome complex (*Nature* 2002, **419:**182-185). They assembled spliceosomes on adenovirus major late pre-mRNA (AdML-M3), which contains three hairpins and can be affinity-purified using a bacteriophage coat protein fused to maltose-binding protein (MS2-MBP fusion protein). Zhou *et al.* demonstrate that the intact complex is highly purified and functional. They examined the protein components by liquid chromatography tandem mass spectrometry (LC-MS/MS) and found around 145 distinct polypeptides; including 88 known splicing-associated proteins. The rest of the proteins had not been previously linked to splicing and include several involved in regulating gene expression, transcription and mRNA export.

References

- 1. Pre-mRNA splicing in the new millennium.
- 2. *Nature*, [http://www.nature.com]