

PUBLISHER CORRECTION

Open Access



# Publisher Correction: Leveraging breeding programs and genomic data in Norway spruce (*Picea abies* L. Karst) for GWAS analysis

Zhi-Qiang Chen<sup>1</sup>, Yanjun Zan<sup>1</sup>, Pascal Milesi<sup>2</sup>, Linghua Zhou<sup>1</sup>, Jun Chen<sup>2,3</sup>, Lili Li<sup>2</sup>, BinBin Cui<sup>4</sup>, Shihui Niu<sup>5</sup>, Johan Westin<sup>6,7</sup>, Bo Karlsson<sup>8</sup>, Maria Rosario García-Gil<sup>1</sup>, Martin Lascoux<sup>2</sup> and Harry X. Wu<sup>1,5,9\*</sup>

The original article can be found online at <https://doi.org/10.1186/s13059-021-02392-1>.

\* Correspondence: [harry.wu@slu.se](mailto:harry.wu@slu.se)

<sup>1</sup>Umeå Plant Science Centre, Department Forest Genetics and Plant Physiology, Swedish University of Agricultural Sciences, SE-90183 Umeå, Sweden

<sup>5</sup>Beijing Advanced Innovation Centre for Tree Breeding by Molecular Design, Beijing Forestry University, Beijing, China

## Publisher Correction to: *Genome Biol* 22, 179 (2021)

<https://doi.org/10.1186/s13059-021-02392-1>

Following publication of the original paper [1], it was reported that Tables 1–3 were published in the incorrect sequence. Tables 1, 2 and 3 were published as Tables 3, 2 and 1 respectively.

Furthermore, there was an error in Fig. 2 and the correct Fig. 2 is supplied below.

The original article [1] has been corrected. The publishers apologize for the error.

### Author details

<sup>1</sup>Umeå Plant Science Centre, Department Forest Genetics and Plant Physiology, Swedish University of Agricultural Sciences, SE-90183 Umeå, Sweden. <sup>2</sup>Program in Plant Ecology and Evolution, Department of Ecology and Genetics, Evolutionary Biology Centre and SciLifeLab, Uppsala University, Uppsala, Sweden. <sup>3</sup>College of Life Sciences, Zhejiang University, Hangzhou 310058, Zhejiang, China. <sup>4</sup>College of Biochemistry and Environmental Engineering, Baoding University, Baoding 071000, Hebei, China. <sup>5</sup>Beijing Advanced Innovation Centre for Tree Breeding by Molecular Design, Beijing Forestry University, Beijing, China. <sup>6</sup>Skogforsk, Box 3, SE-91821 Sävar, Sweden. <sup>7</sup>Unit for Field-Based Forest Research, Swedish University of Agricultural Sciences, SE-90183 Umeå, Sweden. <sup>8</sup>Skogforsk, Ekebo 2250, SE-26890 Svalöv, Sweden. <sup>9</sup>CSIRO National Collection Research Australia, Black Mountain Laboratory, Canberra ACT 2601, Australia.

Published online: 15 July 2021

### Reference

1. Chen ZQ, Zan Y, Milesi P, et al. Leveraging breeding programs and genomic data in Norway spruce (*Picea abies* L. Karst) for GWAS analysis. *Genome Biol.* 2021;22:179 <https://doi.org/10.1186/s13059-021-02392-1>.



© The Author(s). 2021 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

